

Symmetries of the Kadomtsev-Petviashvili Hierarchy

A. Yu. Orlov

Institute of Oceanology, Krasikova 23, Moscow 117218, Russia

P. Winternitz *

Centre de Recherches Mathématiques, Université de Montréal, C.P. 6128 – Succ. Centre-ville, Montréal, Québec H3C 3J7, Canada

The relation between the $\widehat{\mathfrak{sl}}(\infty)$ algebra of flows commuting with the KP hierarchy and the Kac-Moody-Virasoro Lie point symmetries of individual equations is established. This is used to calculate the point symmetries for all equations in the hierarchy.

The Kadomtsev-Petviashvili equation [1] is of considerable physical importance in the theory of wave propagation in such media as shallow water, plasmas and others. It is also the prototype of an integrable equation [2] in $2+1$ dimensions and as such, it has a role to play in conformal quantum field theory and string theories [3]. Like other integrable equations, the KP equation can be viewed as a member of an infinite hierarchy of mutually compatible integrable evolution equations, the KP hierarchy [4,5]. Each equation in the hierarchy involves the same space variables x and y , but a different time variable t_N ($N \geq 3$).

The purpose of this paper is to investigate the symmetries of the KP hierarchy. More specifically we will establish the relation between the $\mathfrak{sl}(\infty)$ symmetries of the hierarchy of equations [4–6] and the Kac-Moody-Virasoro Lie point symmetries of the KP equation itself [7,8]. In the process we will show that each individual equation in the KP hierarchy has a Kac-Moody-Virasoro Lie point symmetry algebra. We shall present the results without proofs, or details. For those we refer to a forthcoming article [9]. In order to present our results, we shall make use of the language of pseudodifferential operators (PDO) in one variable $f(x, \partial)$, with $\partial \equiv \partial/\partial_x$, where f is a function that can be expanded into a Laurent series in ∂ , including an arbitrary number of positive and negative powers [4–6]. We shall use an integral operator

$$K = 1 + \sum_{\ell=1}^{\infty} K_{\ell}(t_1, t_2, t_3, t_4, \dots) \partial^{-\ell} \quad (1)$$

$$t_1 \equiv x, \quad t_2 \equiv y, \quad t_3 = t, \quad \partial \equiv \partial_x.$$

The space of PDO is the direct sum of two subspaces $A = A_+ \dot{+} A_-$, where A_+ includes all nonnegative powers of ∂ , A_- all strictly negative ones.

The equations of the KP hierarchy can now be written in a compact form as

$$\frac{\partial K}{\partial t_n} = -(K \partial^n K^{-1})_- K, \quad (2)$$

where the subscript minus means that only negative powers of ∂ are retained. Putting $\partial_{t_n} \equiv \partial_n$, we can verify that all the flows (2) commute: $[\partial_n, \partial_m]K = 0$. For $n \in \mathbb{Z}^>$ fixed each operator equation (2) represents an infinite coupled set of equations for the coefficients K_{ℓ} in the expansion (1). Various closed subsets of equations can be obtained by fixing n and ℓ . Thus, taking $n \leq 3$, and $\ell = 1, 2, 3$ and appropriately eliminating, we get the potential KP equation for $w \equiv -2K_1$ (with $t_3 = t$)

$$w_{xt} = \frac{1}{4}w_{xxxx} + \frac{3}{2}w_x w_{xx} + \frac{3}{4}w_{yy}. \quad (3)$$

Similarly, for $n = 4$ we get the first higher KP equation

$$w_{xxt_4} = \frac{1}{2}w_{xxxxy} + 3w_{xx}w_{xy} + 2w_x w_{xxy} \\ + w_{xxx}w_y + \frac{1}{2}w_{yyy}. \quad (4)$$

*The research of P.W. was partially supported by research grants from NSERC of Canada of FCAR du Québec. This investigation was conducted during A. Yu. O.'s visit to the CRM, Université de Montréal.

The entire hierarchy can be written as a series of nonlinear and nonlocal evolution equations on $w(x, y, t, t_4, \dots)$, all involving 3 independent variables x, y and t_n (for $n \geq 3$)

$$\frac{\partial w}{\partial t_n} = 2 \operatorname{res}_\partial K \partial^n K^{-1} \quad (5)$$

(where $\operatorname{res}_\partial$ means that we keep only the coefficient of ∂^{-1}).

All equations of the KP hierarchy can be viewed as symmetries of the KP equation. They generate an abelian symmetry algebra and the coefficients of the corresponding vector fields do not depend on the independent variables. Further symmetries exist that do not share the above features: they commute with the KP flows, but not with each other. They can depend explicitly on x, y and t_n . [6], ..., [10] A detailed study of symmetries of the KP hierarchy was performed in Ref. 6. The corresponding equations can be written as

$$\begin{aligned} [\partial_{m,n}, K] &= -(K \hat{x}^n \partial^m K^{-1})_- K \\ \hat{x} &= \sum_{k=1}^{\infty} k t_k \partial^{k-1}, \quad n, m \in \mathbb{Z} \end{aligned} \quad (6)$$

and they correspond to symmetries, since we have $[\partial_{mn}, \partial_k]K = 0$. Their Lie algebra is isomorphic to $\widehat{\mathfrak{sl}}(\infty)$, as can be seen by evaluating the commutator $[\partial_{mn}, \partial_{m'n'}]K$. The operator ∂_{mn} can be viewed as corresponding to differentiation with respect to a new time. The flow of w with respect to this “time” is given by

$$\partial_{mn} w = 2 \operatorname{res}_\partial K \hat{x}^n \partial^m K^{-1}, \quad (7)$$

or equivalently

$$\partial_{mn} w = 2 \operatorname{res}_\lambda \lambda^m \frac{\partial^n \varphi(\lambda)}{\partial \lambda^n} \varphi^*(\lambda), \quad (8)$$

where

$$\varphi(\lambda) = e^\zeta \left(1 + \sum_{k=1}^{\infty} \lambda^k K_n(\vec{t}) \right), \quad \zeta = \sum_{k=1}^{\infty} \lambda^k t_k \quad (9)$$

is the formal Baker-Akhiezer function. [5]

Lie point symmetries, at first glance, do not seem to fit into this scheme. Indeed, e.g. for the KP equation itself, the Lie point symmetries correspond to a transformation $(x, y, t, w) \rightarrow (\tilde{x}, \tilde{y}, \tilde{t}, \tilde{w})$ such that $\tilde{w}(\tilde{x}, \tilde{y}, \tilde{t})$ is a solution, whenever $w(x, y, t)$ is one. The Lie algebra of this Lie group of local point transformations is represented by vector fields of the form

$$\widehat{V} = \xi \partial_x + \eta \partial_y + \tau \partial_t + \phi \partial_w, \quad (10)$$

where ξ, η, τ and ϕ are functions of x, y, t and w . The actual form of these functions is determined using a standard algorithm, [11] having nothing to do with the integrability of the studied equation. Once they are found, a “symmetry”, in the sense of a commuting flow (as in eq. (5), (7), or (8)) is obtained in the form

$$\frac{\partial w}{\partial \lambda} = \xi \frac{\partial w}{\partial x} + \eta \frac{\partial w}{\partial y} + \tau \frac{\partial w}{\partial t} - \phi. \quad (11)$$

Let us now show how the Lie point symmetries fit into the scheme for $\widehat{\mathfrak{sl}}(\infty)$ symmetries (6), ..., (8). We already know [7] that for the KP equation the point symmetry algebra involves arbitrary functions of time. To include these we replace the operators $c_{mn} = \hat{x}^n \partial^m$ by a functions

$$h_{\alpha, N} = \lambda^\alpha h \left(\frac{1}{N \lambda^{N-1}} \frac{\partial}{\partial \lambda} \right) \quad (12)$$

where α and N are integers and h can be expanded into a Laurent series in ∂_E , $E = \lambda^N$. Eq. (8) is then replaced by

$$\begin{aligned} &V^N(\alpha, h)w \\ &= 2 \operatorname{res}_\lambda \lambda^\alpha \left[h \left(\frac{1}{N \lambda^{N-1}} \frac{\partial}{\partial \lambda} \right) \varphi(\lambda) \right] \varphi^*(\lambda) \end{aligned} \quad (13)$$

where V^N is a vector field, acting on w . We shall use eq. (13) to extract all *local* symmetries of the type (13), i.e. those not involving any integrals. We expand the function h in eq. (13), as well as the Baker-Akhiezer functions $\varphi(\lambda)$ and $\varphi^*(\lambda)$. To avoid nonlocal terms in the residue, we keep terms only upto or $O(\lambda^{-N-2})$ in λ (after pulling all derivatives to the right). The result of these operations is

$$\begin{aligned}
V^N(\alpha, h)w = \text{res } \lambda^\alpha & \left\{ h(t_N) + h'(t_N) \left[\frac{2}{N\lambda^{N-2}} y \right. \right. \\
& + \frac{1}{N\lambda^{N-1}} x + \frac{1}{2N\lambda^{N+1}} w \Big] \\
& + \frac{1}{2!} h''(t_N) \left[\frac{1}{N^2\lambda^{2N-4}} y^2 \right. \\
& + \frac{4}{N^2\lambda^{2N-3}} xy + \frac{1}{N^2\lambda^{2N-2}} (x^2 + 4y - 2Ny) \Big] \\
& + \frac{1}{3!} h'''(t_N) \left[\frac{8y^3}{N^3\lambda^{3N-6}} + \frac{12xy^2}{N^3\lambda^{3N-5}} \right] \\
& + \frac{1}{4!} h^{iv}(t_N) \frac{16y^4}{N^4\lambda^{4N-8}} + O(\lambda^{-N-2}) \Big\} \\
& \cdot \left\{ 2 + \frac{w_x}{\lambda^2} + \frac{w_y}{\lambda^3} + \dots + \frac{w_{t_N}}{\lambda^{N+1}} + O(\lambda^{-N-2}) \right\}.
\end{aligned} \tag{14}$$

Fixing $N \geq 3$ in eq. (14) corresponds to choosing an equation in the KP hierarchy. Again, $N = 3$ is the KP equation itself. The value of α corresponds to a specific symmetry. Nonlocal symmetries are avoided by making the restrictions on α . In particular, for $\alpha = N$ we obtain a Virasoro algebra, for $-1 \leq \alpha \leq 2$ we obtain a Kac-Moody type loop subalgebra of the symmetry algebra (in both cases with no central extension). Choosing N and α appropriately, we obtain the Lie point symmetries from eq. (14) in the form of eq. (11), from which we can read off the vector fields (10) directly. Let us run through the individual cases and present the results in terms of vector fields.

$$\begin{aligned}
N = 3, \quad \alpha = 3, 2, 1, 0, -1, \\
T(f) = f\partial_t + \frac{2}{3}yf'\partial_y + \frac{1}{3}\left[xf' + \frac{2}{3}y^2f''\right]\partial_x, \\
\quad - \frac{1}{3}\left[wf' + \frac{1}{3}x^2f'' + \frac{4}{3^2}xy^2f''' + \frac{4}{3^4}y^4f^{iv}\right]\partial_w, \\
Y(g) = g\partial_y + \frac{2}{3}yg'\partial_x - \frac{4}{9}y\left(xg'' + \frac{2}{9}y^2g'''\right)\partial_w, \\
X(h) = h\partial_x - \frac{2}{3}\left(xh' + \frac{2}{3}y^2h''\right)\partial_w, \\
W(k) = ky\partial_w, \quad U(\ell) = \ell\partial_w.
\end{aligned} \tag{15}$$

$$\begin{aligned}
N = 4, \quad \alpha = 4, 2, 1, -1 \\
T(f) = f\partial_t + \frac{1}{4}f'(x\partial_x + 2y\partial_y) \\
\quad - \frac{1}{4}(wf' + xyf'')\partial_w, \\
Y(g) = g\partial_y - \frac{1}{2}xg'\partial_w, \\
X(h) = h\partial_x - yh'\partial_w, \quad W(k) = k\partial_w.
\end{aligned} \tag{16}$$

$$\begin{aligned}
N = 5, \quad \alpha = 5, 2, 1, -1, \\
T(f) = f\partial_t + \frac{1}{5}f'(x\partial_x + 2y\partial_y) \\
\quad - \left(\frac{1}{5}wf' + \frac{4}{25}y^2f''\right)\partial_w, \\
Y(g) = g\partial_y - \frac{4}{5}g'y\partial_w, \quad X(h) = h\partial_x, \\
W(k) = k\partial_w,
\end{aligned} \tag{17}$$

$$\begin{aligned}
N \geq 6, \quad \alpha = 6, 2, 1, -1, \\
T(f) = f\partial_t + \frac{1}{N}f'(x\partial_x + 2y\partial_y - w\partial_w), \\
Y(g) = g\partial_y, \quad X(h) = h\partial_x, \\
W(k) = k\partial_w.
\end{aligned} \tag{18}$$

In eq. (15), ..., (18) all of the functions f, g, h, k and ℓ depend on the corresponding time t_N and are otherwise arbitrary.

Eq. (15) for the potential KP agrees with ref. [9], the cases $N \geq 4$ are new. The first field $T(f)$ in each case corresponds to the Virasoro algebra, the remaining ones to loop algebras, based on solvable Lie algebras. Notice that the Lie point symmetries go far beyond the obvious “physical” symmetries. Indeed time translations, and dilations are obtained from the Virasoro algebra $T(f)$ as $T(1)$ and $T(t)$, respectively. In general, $T(f)$ generates arbitrary reparametrizations of time. Similarly, $X(1)$ is an x -translation, $X(t)$ a Galilei transformation; $Y(1)$ a y -translation, $Y(t)$ a Galilei transformation for $N \geq 4$. For $N = 3$ it is a Galilei transformation combined with a rotation. Finally, $W(k)$ and $U(\ell)$ generate gauge transformations.

We mention that all known integrable equations in $2 + 1$ dimensions have Kac-Moody-Virasoro Lie point symmetry algebras (Davey-Stewartson equation, 3-wave equations, two dimensional Toda lattice, etc.). [12] The results presented here show that this is no coincidence and give credence to the conjecture that the occurrence of Kac-Moody-Virasoro symmetries can be used as an integrability indicator for PDEs in 3-dimensions of space-time.

- [1] V. V. Kadomtsev and V. I. Petviashvili, Soviet Phys. Dokl. **15**, 539 (1970).
- [2] M. J. Ablowitz and P. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- [3] D. Gross and A. Migdal, Phys. Rev. Lett. **64**, 127 (1990); E. Witten, Nuclear Phys. B **340**, 281 (1990); F. Yu and Y. S. Wu, J. Math. Phys. **34**, 5851 (1993); A. Gerasimov, A. Marshakov, A. Mironov, A. Morozov, and A. Orlov, Nuclear Phys. B **357**, 565 (1991).
- [4] M. Jimbo and T. Miwa, Publ. RIMS Kyoto University **19**, 943 (1983).
- [5] L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, World Scientific, Singapore, 1991.
- [6] A. Yu. Orlov and E. I. Schulman, Lett. Math. Phys. **12**, 171 (1986), Novosibirsk preprint, 217 (1984), 277 (1985).
- [7] D. David, N. Kamran, D. Levi, and P. Winternitz, Phys. Rev. Lett. **55**, 2111 (1985), J. Math. Phys. **27**, 1225 (1986); D. David, D. Levi and P. Winternitz, Phys. Lett. A **118**, 390 (1986).
- [8] B. Dorizzi, B. Grammaticos, A. Ramani, and P. Winternitz, J. Math. Phys. **27** 2848 (1986); J. Rubin and P. Winternitz, J. Math. Phys. **31** 2085 (1990).
- [9] A. Yu. Orlov and P. Winternitz, Preprint CRM-1936, Montréal, 1994, to be published.
- [10] F. Schwarz, J. Phys. Soc. Japan. **51**, 2387 (1982); E. Infeld and P. Frycz, Acta Phys. Polon. B **14**, 129 (1983); H. H. Chen, Y. C. Lee and J. E. Lin, Physica **D9**, 439 (1983).
- [11] P. J. Olver, *Application of Lie Groups to Differential Equations*, Springer, Berlin, 1986.
- [12] B. Champagne and P. Winternitz, J. Math. Phys. **29**, 1 (1988); L. Martina and P. Winternitz, Ann. Physics (N.Y.) **196**, 231 (1989); D. Levi and P. Winternitz, J. Math. Phys. **34**, 3713 (1993).